arXiv:1508.00631v1 [astro-ph.CO] 4 Aug 2015

The fate of a Universe driven by a linear potential

Ricardo Z. Ferreira*
CP^3-Origins, Center for Cosmology and Particle Physics Phenomenology
University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark

Pedro P. Avelino†
Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto,
CAUP, Rua das Estrelas, PT4150-762 Porto, Portugal
Centro de Astrofísica da Universidade do Porto,
Rua das Estrelas, , PT4150-762 Porto, Portugal and
Departamento de Física e Astronomia, Faculdade de Ciências,
Universidade do Porto, Rua do Campo Alegre 687, PT4169-007 Porto, Portugal

We study the fate of our Universe assuming that the present accelerated stage is due to a scalar field in a linear potential. Such a Universe would bounce and collapse in the future. We solve numerically and analytically the equations of motion for the scalar field and the scale factor. In particular, we relate the duration of the accelerated stage, the bounce and the collapse with the mass of the field and, thus, with the current value of equation of state w. We obtain an expression which predicts the age of the Universe for a given $w+1$. The present constraints on w imply that the Universe will not collapse in the next 56 billion years. Moreover, a cosmological solution to the coincidence problem favors a significant deviation of w from -1 such that the Universe collapses in the not too distant future.

I. INTRODUCTION

The Cosmological Constant (CC) problem [1–4] is a shadow in modern Physics. Such a discrepancy between predictions and observations has been, in the history of Science, a sign of a paradigmatic change of our understanding of the Universe. It has been tackled in the last decades in several different ways, all of them with a common negative outcome (see [5] for a recent review). Nevertheless, given the relevance of the problem one should keep an open mind for new proposals like the recent one dubbed vacuum energy sequestering mechanism (VES) [6, 7].

Here we address the *new* version of the problem known as the coincidence problem. It consists on the coincidence between the epoch we live in and the beginning of an epoch where the Universe is dominated by what looks like a CC. Although the present data is consistent with a pure CC ($w = -1.019^{+0.075}_{-0.080}$ (95% CL) [8]) there are several different ways to drive the present accelerated stage [9]. One of the most interesting alternative ideas relies on a scalar (quintessence) field slowly-rolling its potential until the present time, when it starts dominating the energy content of the Universe driving a period of accelerated expansion [10] (see also [11] for a more recent analysis).

In particular, a subclass of these quintessence models has received much attention due to its technical naturalness which protects a small mass for the field [12–14], needed to explain the observations. That subclass consists of scenarios where the scalar field has a shift symmetry only broken by non-perturbative effects. Specific realizations include cosine but also some monomial potentials [15–17] for the field. Here we will focus on the simple case of a linear potential. The use of a linear potential to explain the current dark energy stage has been a recurrent idea in the literature [18–23]. Apart from the virtues mentioned above, the case of the linear potential is also appealing because its dynamics is quite insensitive to the initial conditions. A light field is essentially frozen in its potential, dynamically this stage is quickly achieved independently of the initial conditions, until it dominates the energy content of the Universe and drives a period of accelerated expansion. However, this period does not last forever. At some point in the future the slow-roll conditions are broken and the dark energy stage ends. At that time, the field quickly rolls down its potential to negative values such that the energy density of the Universe goes to zero, the Universe bounces and a collapsing stage starts dominated by the kinetic energy of the field [20]. In this work we perform a detailed analytical and numerical analysis of the dynamics. We discuss the coincidence problem in this context by establishing a relation between the total age of the Universe and the value of the equation of state w today. A future observation of $w > -1$ would then indicate the time until the *doomsday*.

The paper is organized as follows. In section II we study the dynamics of a linear potential in an expanding Universe. We solve numerically and analytically the 3 relevant stages: matter domination, accelerated expansion and the collapse. Then, assuming that the linear potential drives the present stage of acceleration we derive relations between the equation of state and the time until the collapse. In section III we conclude.

We assume a FRW metric in cosmic time (t) with a

---

* ferreira@cp3.dias.sdu.dk
† pedro.avelino@astro.up.pt

small positive curvature (k) which will not play any important role in the dynamics.

II. LINEAR POTENTIAL AND DARK ENERGY

A. Dynamics

We consider a scalar field ϕ with a linear potential

$$V(\phi) = m^3\phi, \tag{1}$$

where m is the mass of the field. The equation of motion for a homogeneous scalar field in a linear potential is given by

$$\ddot{\phi} + 3H\dot{\phi} + m^3 = 0, \tag{2}$$

where $H = \dot{a}/a$ is the Hubble rate, a is the scale factor and the dot means a derivative with respect to cosmic time.

1. Radiation/Matter stage

We start the analysis at a period where the Universe is dominated by some perfect fluid with equation of state w, hence, the scale factor evolves in time as $a = a_0(t/t_0)^p$ where $p = 2/(3(1+w)) > 1/3$. For such a background evolution the scalar field behaves as

$$\phi(t) = C_1 + \frac{C_2}{(1-3p)t^{3p-1}} - \frac{m^3t^2}{2(3p+1)}, \tag{3}$$

where C_1, C_2 are two integration constants. For reasonable initial conditions one expects the second term in the right hand side to be diluted and so, at sufficient late times, we effectively have

$$\phi = \phi_i - \frac{m^3t^2}{2(3p+1)}. \tag{4}$$

As one would expect, if the field is light, $H = p/t \gg m$, the second term on the rhs of Eq. (4) is small and so the field remains effectively frozen in its potential. Even for masses close to the maximal value allowed by observations the value of ϕ today would only differ $\simeq 17\%$ from ϕ_i.

2. Accelerated stage

While the Universe expands, H decreases and we reach a time at which ϕ dominates the energy content of the Universe: $\rho_{\rm total} = 3H_{\rm DE}^2M_p^2 \simeq \rho_\phi \simeq m^3\phi_i$. If $\phi > 0$ the field drives a period of accelerated expansion. Depending on the mass of the field, this period can be well described by the slow-roll approximation which requires the slow-roll parameters $\epsilon \equiv \dot{\phi}^2/(2H^2M_p^2)$ and $\delta = \ddot{\phi}/(H\dot{\phi})$ to be $\ll 1$ (in absolute value). The slow-roll condition for ϵ requires:

$$\left(\frac{m^3t_{\rm DE}}{3p+1}\right)^2 \frac{1}{2H_{\rm DE}^2M_p^2} \ll 1, \tag{5}$$

where M_p is the reduced Planck mass. On the other hand, the condition for δ is

$$(H_{\rm DE}t_{\rm DE})^{-1} \ll 1. \tag{6}$$

If these conditions are satisfied then the dynamics of ϕ is no longer given by Eq. (4). Instead, using the slow-roll approximation to neglect the second derivative of ϕ in Eq. (2) and assuming $\rho_{\rm total} = m^3\phi_i$, we have

$$\phi(t) \simeq \left[\phi_i^{3/2} - \frac{\sqrt{3m^3}}{2}(t - t_{\rm DE})M_p\right]^{2/3}. \tag{7}$$

The accelerated period will end at a time t_s when the slow-roll conditions are violated $\epsilon(t_s) \simeq 1$. Using Eq. (7) we can estimate the duration of the accelerated stage to be

$$t_s - t_{\rm DE} = \frac{2}{\sqrt{3m^3}M_p}\left[\phi_i^{3/2} - \left(\frac{M_p}{\sqrt{2}}\right)^{3/2}\right]. \tag{8}$$

Note that this analysis is only valid if $\phi_i \gtrsim M_p/\sqrt{2}$. This can be seen as a consistency condition to have slow-roll in the first place.

For times close to the present time (t_0) then $H^{-1} \simeq t$ and the first slow-roll condition (ϵ) requires $m^3 \ll H_{\rm DE}^2M_p$. On the other hand the second slow-roll parameter (δ) is never $\ll 1$ around the present epoch. Therefore, the dynamics is only accurately described by slow-roll at a time $t_{\rm DE} > t_0$. In particular, for the largest masses allowed by the observations, the slow-roll condition for δ is never strictly satisfied, even at late times. Nevertheless, we will assume the slow-roll to be is valid as long as $\epsilon, \delta < 1$ (which is typically satisfied even before today). Naturally, this assumption will insert a mild error in the analytical results when comparing to the numerical results. In particular, the fact that the slow-roll conditions are not strictly satisfied today implies an error in estimating $\dot{\phi}(t_0)$ which, as we will explain in more detail later, affects the computation of the equation of state $w(t_0)$.

The initial condition for ϕ is not a free parameter if we want to explain the present period of acceleration. In that case[1]

$$\phi_i m^3 = \Omega_{\rm DE}\rho_0, \tag{9}$$

where $\rho_0 = 3H_0^2M_p^2$ is the energy density today and $\Omega_{\rm DE} \simeq 0.69$ [8] is the dark energy density parameter today.

---

[1] To be more precise $\phi_i m^3 + \dot{\phi}^2/2 = \Omega_{\rm DE}\rho_0$. However, in the cases allowed by observations this should give a small correction.



FIG. 1. The total energy density of the Universe (black), the energy density in matter (blue) and the energy density in the field ϕ (red, dashed is analytical) from an early time until the collapse. We choose $m^3 = 0.538\, H_0^2 M_p$ ($w = 0.01$), $\Omega_{\rm DE} = 0.69$ and no curvature. The energy densities are normalized by ρ_0 and the times by t_0. In green and purple is plotted the slow-roll parameters ϵ and δ, respectively.

In Fig. (1) we show the dynamics of the Universe described in the previous paragraphs containing only matter and a scalar field (ϕ). We fixed $\phi_i = \phi(t_0)$ through Eq. (9) and choose $m^3 = 0.538\, H_0^2 M_p$ ($w+1 = 0.01$). The value of $\dot\phi$ is fixed by assuming that Eq. (4) is satisfied well inside the matter dominated era. We can verify that the analytical solution for ϕ in Eq. (7) is accurate until the time the slow-roll conditions are violated. Moreover, for such a value of the mass Eq. (8) predicts the duration of the dark energy stage to be $t_s - t_{\rm DE} \simeq 11.5\, H_0^{-1}$ which is in reasonable agreement with the numerical result of $12.9\, t_0$. The difference corresponds to small deviations between ϕ_i and $\phi(t_{\rm DE})$, and the fact that our conditions for slow-roll requirements are not exact. We verified that the error decreases for small masses while for the largest masses compatible with observations ($w+1 \simeq 0.1$), where the slow-roll is never strictly achieved, the error is $\simeq 50\%$.

3. *Bounce and collapse*

At the time the slow-roll conditions are violated the field starts rolling down the potential until it reaches a point where $\phi < 0$ and $m^3\phi + \dot\phi^2/2 = 0$. Thus, the total energy of the Universe goes to zero ($H \to 0$) and a bounce occurs. The influence of other components like matter, radiation or curvature is, at this point, very much diluted and so it will no interfere in the dynamics.

During this particular time window, since the slow-roll condition is violated (t_s) until the bounce (t_b), it is difficult to have an analytical understanding of the dynamics. However, we can still get an analytical insight by noting that the kinetic $K(\phi)$ energy of ϕ is the dominant contribution for most of the time. Therefore, the equation of motion for ϕ will be approximately given by

$$\ddot\phi + \sqrt{\frac{3}{2}}\frac{\left|\dot\phi\right|\dot\phi}{M_p} + m^3 = 0. \tag{10}$$

As $\dot\phi < 0$ the solution for this equation, after matching with Eq. (7) at t_s, yields

$$\frac{\phi(t)}{M_p} = \frac{1}{\sqrt{2}} + \frac{\log\left(\frac{1}{2}(3+\sqrt{3})\right)}{\sqrt{6}} - \sqrt{\frac{2}{3}}bM_p \times \\ \times \log\left[\cosh\left[b(t-t_s) + \operatorname{arctanh}\left(3^{-1/4}\right)\right]\right] \tag{11}$$

where $b = (3/2)^{1/4}\sqrt{m^3/M_p}$, while $\dot\phi$ is given by

$$\dot\phi(t) = -\sqrt{\frac{2}{3}}bM_p \tanh\left[b(t-t_s) + \operatorname{arccoth}\left(3^{1/4}\right)\right] \tag{12}$$

The previous equation saturates before the bounce to $\dot\phi(t) \simeq -\sqrt{2/3}\, bM_p$. The bounce itself will happen when $m^3\phi + \dot\phi^2/2 = 0$. Using the saturated value for $\dot\phi$ we get $\phi(t_b) \simeq -M_p/\sqrt{6}$. Then, by inserting this result into Eq. (11) we find the time between t_s and t_b to be

$$t_b - t_s \simeq 1.34\sqrt{\frac{M_p}{m^3}}. \tag{13}$$

When $H = 0$ the acceleration is negative so the Universe starts to collapse dominated again by the kinetic energy of ϕ, in order to keep $\rho_{\rm total} > 0$, while the field continues to roll down the potential. Thus, for $t > t_b$ the equation of motion for ϕ becomes

$$\ddot\phi - \sqrt{\frac{3}{2}}\frac{\dot\phi|\dot\phi|}{M_p} + m^3 = 0, \tag{14}$$

which is similar to Eq. (10) apart from the fact that $H < 0$ during the collapse. As $|\dot\phi| = -\dot\phi$ the solution for $\dot\phi$ is now given by

$$\dot\phi(t) = -\sqrt{\frac{2}{3}}bM_p \tan\left[b(t-t_f) + \frac{\pi}{2}\right], \tag{15}$$

where t_f is an integration constant. Note that the solution is only valid for $0 < b(t_f - t) < \pi/2$. When $t - t_f = 0$, $\dot\phi$ diverges. This divergence corresponds to the collapse of the Universe as we can verify by solving the Friedmann equation for the scale factor

$$a(t) = C_4\left[\cos\left(b(t-t_f) + \frac{\pi}{2}\right)\right]^{1/3}, \tag{16}$$

where C_4 is another integration constant. When $t - t_f \to 0$ the scale factor tends to zero. Therefore, we can identify t_f with the total age of the Universe. Close to t_f



FIG. 2. The total energy density of the Universe (black) and the kinetic $K(\phi)$ (red) and potential $V(\phi)$ (blue) energy of ϕ for $m^3 = 0.538\, H_0^2 M_p$ ($w = 0.01$). The blue dashed line uses the analytical approximation for ϕ before the bounce Eq. (11) while the the red dashed line uses the analytical solution for $\dot{\phi}$ during the collapse Eq. (18). The energy densities are normalized by ρ_0.

the solutions can be expanded to leading order in $t_f - t$ giving

$$a(t) = a_b \left(\frac{t_f - t}{t_f - t_b}\right)^{1/3}, \tag{17}$$

where we have fixed $C_4 = a_b/(b(t_f - t_b))^{1/3}$, and

$$\dot{\phi} = -\sqrt{\frac{2}{3}}\frac{M_p}{t_f - t}. \tag{18}$$

These results are in agreement with what we would expect from a Universe dominated by a kinetic stage ($w \simeq 1$) [20]. The final relevant quantity left to determine is the total age of the Universe t_f. In order to compute it we match Eqs. (12) and (18) at the bounce. Using the saturated value for $\dot{\phi}$ in Eq. (12) we simply get

$$t_f - t_b \simeq b^{-1} = \left(\sqrt{\frac{2}{3}}\frac{M_p}{m^3}\right)^{1/2}. \tag{19}$$

In sum, both time windows, from the end of the dark energy stage to the bounce and from the bounce to the collapse are $\mathcal{O}(\sqrt{M_p/m^3})$.

In Fig. (2) we present the dynamics of the Universe, with the same conditions as in Fig. (1), zoomed in the region from the end of the dark energy stage (t_s) to the complete collapse (t_f), where the energy density diverges. For the set of parameters chosen, the slow-roll conditions are violated at $t_s \simeq 13.64\, t_0$, the bounce occurs at $t_b \simeq 15.82\, t_0$ while the collapse happens at $t_f \simeq 17.07\, t_0$. In the region before the bounce we can verify that Eq. (11) is a reasonable approximation for ϕ by comparing the blue lines. Our expression is fully analytical and therefore uses the analytical approximations for ϕ and $\dot{\phi}$ computed at the end of slow-roll. That introduces a small error in this approximation. In Eq. (13) we estimated the time window from t_s to t_b as a function of the mass of the field. For the mass used here it yields $t_b - t_s = 1.91\, t_0$ which is close to the numerical value of $2.17\, t_0$.

Regarding the collapsing stage, the total energy in the collapse is basically given by the kinetic energy of ϕ, as expected. Furthermore, we can verify that the solution for $\dot{\phi}$ obtained in Eq. (18) is a good approximation during the collapse. Using Eq. (19) we estimate $t_f - t_b = 1.29\, t_0$ which is in very good agreement with the numerical value of $1.25\, t_0$.

We verified for other values of the mass that our analytical estimations for $t_f - t_b$ is accurate with an error $< 3\%$ while the estimation of $t_b - t_s$ is also good with an error $< 15\%$. Moreover, we have also checked that the addition of curvature consistent with observations does not affect the dynamics.

B. Predictions

The previous analysis allowed us to derived relations between the mass of the field, the duration of the accelerated stage, the bounce and the collapse. Recall that we assumed $t_{\rm DE}$ to be the time at which the slow-roll parameters are < 1 and so $t_{\rm DE} < t_0$. For the range of masses analyzed $t_{\rm DE} \simeq 0.72\, t_0$. Therefore, using Eqs. (8), (13) and (19) we can write the total age of the Universe only as a function of the mass of the field as

$$t_f \simeq \frac{6\Omega_{\rm DE}^{3/2} H_0^3 M_p^2}{m^6} + 1.56\sqrt{\frac{M_p}{m^3}} + 0.72 t_0. \tag{20}$$

It is now interesting and useful to rewrite these relations in terms of the observed equation of state parameter w. Recall that the equation of state is measured by observations to be very close to a pure CC: $w = -1.019^{+0.075}_{-0.080}$ (95% CL) [8]. In order to make predictions of what a given value of w would tell us about the fate of the Universe we use the fact that during the dark energy stage

$$w = \frac{K(\phi) - V(\phi)}{K(\phi) + V(\phi)} \simeq -1 + \frac{2K(\phi)}{V(\phi)}. \tag{21}$$

Therefore, using the slow-roll conditions, the equation of state today is given by

$$w + 1 = \frac{\Omega_{\rm DE} M_p^2}{3\phi_i^2} = \frac{m^6 M_p^2}{3\Omega_{\rm DE}\rho_0^2}, \tag{22}$$

where we have used Eqs. (7) and (9). However, as we explained in the previous section, the present epoch is a transient regime where slow-roll is not a very accurate



FIG. 3. The coincidence problem, defined as the ratio between the total age of the Universe over the time the Universe has been accelerating so far, as a function of the equation of state today. The blue line is the analytical result while the black is the numerical.

approximation. Both Eq. (4) and Eq. (7) overestimate $\dot{\phi}(t_0)$. A possible improvement would be to solve the equations assuming a Universe dominated by CC and matter until today and then solve the equation for the scalar field in such background. We take a more simplistic approach by noting that the previous expression overestimates the value of $w+1$ by a factor of $\gamma \simeq 1.5$. This is true for a wide range of $w+1$, which again suggests that an analytical improvement might be easily achieved. In what follows we take this factor γ into consideration. Therefore, combining Eqs. (20) and (22) gives

$$t_f = \frac{\sqrt{\Omega_{\rm DE}}}{H_0(w+1)\gamma}\left[\frac{2}{9} + \frac{0.68}{\Omega_{\rm DE}^{5/4}}\gamma^{3/4}(w+1)^{3/4}\right] + 0.72t_0. \tag{23}$$

For $w+1 \ll 1$ the age of the Universe is inversely proportional to $w+1$. The present observational constraint of $w+1 \lesssim \mathcal{O}(0.1)$ implies that the collapse of the Universe, an so the *doomsday*, will not happen in the next 56 billion years. The previous equation underestimates the age of the Universe for large masses with a maximal error of 16% which decreases for small masses, as explained in the previous section.

We are now in position to properly analyze the coincidence problem in the context of the linear potential driving dark energy. The fact that the linear potential triggers a collapse of the Universe ameliorates per se the coincidence problem as the Universe is no longer eternal. We can quantify this improvement by defining the ratio between the time elapsed since the Universe started accelerating ($t_{\rm acel}$) until today, with the total age of the Universe

$$f = \frac{t_0 - t_{\rm acel}}{t_f}. \tag{24}$$

In the range of $w+1$ considered $t_0 - t_{\rm acel} \simeq 0.45\,t_0$ although the precise value varies with the mass of the field. In Fig. (3) we present this quantity evaluated both analytically, through Eq. (23), and numerically. The numerical results are in agreement with those ones obtained in [20]. Our numerical expression is very accurate for small masses although it underestimates the age of the Universe for larger masses. In the context of the coincidence problem these results show that a valid solution requires a dynamical field with an equation of state not to far from the current bounds and, therefore, possibly observed with the forthcoming experiments. In particular, if $w+1$ is close to the maximal value allowed by observations the coincidence would only be at the level of 1/10 while $w+1 \simeq 0.01$ would represent a 1/30 level of coincidence.

III. CONCLUSION

In this work we studied in detail the fate of the Universe assuming that the current accelerated stage is driven by a scalar field in a linear potential. The case of a linear potential is of great interest because of its predictive dynamics, quite insensitive to the initial conditions, but also because a small mass of the field can be made technically natural. Generically, a scalar field in a linear potential and in an expanding Universe will, at some point, dominate the energy density and drive a period of dark energy followed by a bounce and, finally, a crunch. We have derived analytical expressions for the scalar field and the scale factor during these epochs. This allowed us to estimate the time duration of the several stages as a function of the mass of the field. The analytical expressions are in good agreement with the numerical results, in particular, the collapsing time estimation in Eq. (19). The accuracy of our expressions increases for smaller masses while for masses close to the largest values allowed by observations there is a larger error because the slow-roll dynamics is no longer the best description for the scalar field. From the expressions derived in the three relevant stages we derived our main result, Eq. (23), where we predict the age of the universe for a given observed value of the equation of state w. Our expression is accurate for smaller masses although it underestimates the age of the Universe for the largest masses compatible with observations. The present observational constraints of $w+1 \lesssim 0.1$ enforces that the *doomsday* does not happen in the next 56 billion years. Finally, we also compute how the linear potential addresses the coincidence problem by computing the ratio between the time the Universe was accelerating and its total age (see Fig. (3)). We find a 1/10 level of coincidence for $w+1 \simeq 0.1$, which is very reasonable, and 1/30 for $w+1 \simeq 0.01$, which although acceptable is border line to what we would consider a solution to the coincidence problem. Therefore, a solution to the coincidence problem prefers a more dynamical dark energy stage with a value of $w + 1$ possibly observed by future experiments.

[1] Y. Zeldovich, JETP Lett. **6**, 316 (1967).

[2] Y. Zel'dovich, A. Krasinski, and Y. Zeldovich, Sov.Phys.Usp. **11**, 381 (1968).

[3] F. Wilczek, Phys.Rept. **104**, 143 (1984).

[4] S. Weinberg, Rev.Mod.Phys. **61**, 1 (1989).

[5] C. Burgess, (2013), arXiv:1309.4133 [hep-th].

[6] N. Kaloper and A. Padilla, Phys.Rev.Lett. **112**, 091304 (2014), arXiv:1309.6562 [hep-th].

[7] N. Kaloper and A. Padilla, Phys.Rev. **D90**, 084023 (2014), arXiv:1406.0711 [hep-th].

[8] P. Ade *et al.* (Planck), (2015), arXiv:1502.01589 [astro-ph.CO].

[9] E. J. Copeland, M. Sami, and S. Tsujikawa, Int.J.Mod.Phys. **D15**, 1753 (2006), arXiv:hep-th/0603057 [hep-th].

[10] B. Ratra and P. Peebles, Phys.Rev. **D37**, 3406 (1988).

[11] E. V. Linder, Phys.Rev. **D91**, 063006 (2015), arXiv:1501.01634 [astro-ph.CO].

[12] J. A. Frieman, C. T. Hill, and R. Watkins, Phys.Rev. **D46**, 1226 (1992).

[13] C. T. Hill, D. N. Schramm, and J. N. Fry, Comments Nucl.Part.Phys. **19**, 25 (1989).

[14] J. A. Frieman, C. T. Hill, A. Stebbins, and I. Waga, Phys.Rev.Lett. **75**, 2077 (1995), arXiv:astro-ph/9505060 [astro-ph].

[15] L. McAllister, E. Silverstein, and A. Westphal, Phys.Rev. **D82**, 046003 (2010), arXiv:0808.0706 [hep-th].

[16] N. Kaloper and L. Sorbo, Phys.Rev.Lett. **102**, 121301 (2009), arXiv:0811.1989 [hep-th].

[17] N. Kaloper, A. Lawrence, and L. Sorbo, JCAP **1103**, 023 (2011), arXiv:1101.0026 [hep-th].

[18] R. Kallosh, J. Kratochvil, A. D. Linde, E. V. Linder, and M. Shmakova, JCAP **0310**, 015 (2003), arXiv:astro-ph/0307185 [astro-ph].

[19] P. Avelino, C. Martins, and J. Oliveira, Phys.Rev. **D70**, 083506 (2004), arXiv:astro-ph/0402379 [astro-ph].

[20] P. P. Avelino, Phys.Lett. **B611**, 15 (2005), arXiv:astro-ph/0411033 [astro-ph].

[21] E. V. Linder, Gen.Rel.Grav. **40**, 329 (2008), arXiv:0704.2064 [astro-ph].

[22] P. Avelino, Phys.Rev. **D90**, 103523 (2014), arXiv:1410.4555 [gr-qc].

[23] A. Barreira and P. Avelino, Phys.Rev. **D83**, 103001 (2011), arXiv:1103.2401 [astro-ph.CO].